



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

In this instrument the alternations in brightness at any point in the field when the slide is moved are beats due to the Doppler effect just as truly as are those heard in the second form of Koenig's experiment.

ALBERT B. PORTER.

CHICAGO,
January 14, 1905.

NOTE ON THE BROAD WHITE FISH.

In the *Proceedings* of the American Philosophical Society of Philadelphia, XLIII., 1904, p. 451, plates VIII. and IX., I have wrongly identified the broad white fish, or *Coregonus kennicotti* Jordan and Gilbert, as the humpback, or *Coregonus nelsonii* Bean. My error was due largely to lack of material, ignorance of the species from autopsy, and the fact, as I have since discovered, that *C. nelsonii* does not always exhibit the well-developed hump like that of the type. Possibly when the Siberian forms are carefully studied the nomenclatures of these fishes will be more stable.

HENRY W. FOWLER.

ACADEMY OF NATURAL SCIENCES,
PHILADELPHIA, February 5, 1905.

RECENT ZOOPALEONTOLOGY.*

DURING the past thirteen years great advances have been made in our knowledge of the ancient mammalian life of North America, especially through the explorations in the Rocky Mountain region carried on by the Carnegie, Field Columbian and American Natural History Museums. The long Tertiary period has been clearly subdivided into a series of stages and substages. This enables paleontologists to record more accurately than ever before the time of arrival and departure of the larger and smaller quadrupeds from North and South America, Asia, Europe, Africa, and to determine more precisely when the connection of North and South America was interrupted by a gulf flowing between the Atlantic and Pacific Oceans, and when the connection was again made by the elevation

* Abstract of a lecture delivered by Professor Osborn before the Society of Naturalists at the Philadelphia meeting.

of the Isthmus of Panama; this demonstrates also that a very much closer connection existed between the animal life of Europe and of North America through continuous intermigration over the broad land area now submerged beneath the Behring Straits. A series of six world maps prepared by Dr. W. D. Matthew clearly exhibit this submergence and emergence of the isthmuses between these great continents.

Of especial interest is the recent discovery by the Geological Survey of Egypt that the whole race of mastodons and elephants originated in Africa, entered Europe in the middle of the Tertiary and soon afterward found their way into North America and somewhat later into South America. We have now been able to fix very positively the date of actual arrival of these animals in North America. It appears probable that successive waves of migrations of European and Asiatic species of elephants and mammoths came to this country. In the meantime there survived here from one of the earliest African migrants the eastern American forest mastodon which lived until comparatively recent times.

The theory of multiple races or polyphyletic evolution not only of elephants but of horses, rhinoceroses, camels and titanotheres appears to be clearly established through these recent discoveries. It was formerly believed, for example, that the modern horse had a single line of ancestors extending back into the Eocene period; now it appears that in North America there were always four to six entirely different varieties of the horse family living contemporaneously, including slow-moving, forest-living horses with broader feet, and very swift plains-living horses with narrow feet fashioned more like the deer. Intermediate between these arose the variety which survived and gave rise to the true modern horse. Furthermore, it appears that the modern horses separated from the asses and zebras at a much more remote period than has been generally supposed, and we are now endeavoring to ascertain accurately when this separation occurred.

The same discovery of multiple races has been made among the rhinoceroses. In Eu-